

Killing niche competitors by remote-control bacteriophage induction

Laura Selva^a, David Viana^a, Gili Regev-Yochay^b, Krzysztof Trzcinski^b, Juan Manuel Corpa^a, Íñigo Lasa^c, Richard P. Novick^d, and José R. Penadés^{a,e,1}

^aUniversidad Cardenal Herrera-CEU, 46113 Moncada, Valencia, Spain; ^bDepartments of Epidemiology and Immunology and Infectious Diseases, Harvard School of Public Health, Boston; ^cInstituto de Agrobiotecnología, CSIC-Universidad Pública de Navarra, 31006 Pamplona, Navarra, Spain; ^dSkirball Institute, New York University Medical Center, 540 First Avenue, New York, NY 10016; and ^eCentro de Investigación y Tecnología Animal, Instituto Valenciano de Investigaciones Agrarias (CITA-IVIA), Apdo. 187, 12.400 Segorbe, Castellón, Spain

Edited by Sankar Adhya, National Institutes of Health, Bethesda, MD, and approved December 3, 2008 (received for review September 26, 2008)

A surprising example of interspecies competition is the production by certain bacteria of hydrogen peroxide at concentrations that are lethal for others. A case in point is the displacement of *Staphylococcus aureus* by *Streptococcus pneumoniae* in the nasopharynx, which is of considerable clinical significance. How it is accomplished, however, has been a great mystery, because H₂O₂ is a very well known disinfectant whose lethality is largely due to the production of hyperoxides through the abiological Fenton reaction. In this report, we have solved the mystery by showing that H₂O₂ at the concentrations typically produced by pneumococci kills lysogenic but not nonlysogenic staphylococci by inducing the SOS response. The SOS response, a stress response to DNA damage, not only invokes DNA repair mechanisms but also induces resident prophages, and the resulting lysis is responsible for H₂O₂ lethality. Because the vast majority of *S. aureus* strains are lysogenic, the production of H₂O₂ is a very widely effective antistaphylococcal strategy. Pneumococci, however, which are also commonly lysogenic and undergo SOS induction in response to DNA-damaging agents such as mitomycin C, are not SOS-induced on exposure to H₂O₂. This is apparently because they are resistant to the DNA-damaging effects of the Fenton reaction. The production of an SOS-inducing signal to activate prophages in neighboring organisms is thus a rather unique competitive strategy, which we suggest may be in widespread use for bacterial interference. However, this strategy has as a by-product the release of active phage, which can potentially spread mobile genetic elements carrying virulence genes.

hydrogen peroxide | SOS response | *Staphylococcus aureus* | *Streptococcus pneumoniae* | bacterial interference

The interactions among bacteria living communally are highly complex and extremely interesting—illuminating, as they do, a long-ignored but nevertheless critical aspect of microbial biology. One can readily envision interactions such as direct competition for scarce nutrients, mutual cooperation for the conversion of substrates to utilizable metabolites, “borrowing” of quorum-sensing signals, DNA transfer, biofilm formation and maintenance, and interference or inhibition mediated by antibacterial products, including bacteriocins, antibiotics, and low-molecular-weight toxic compounds such as H₂O₂ (1). In this article, we consider a specific case of H₂O₂-mediated bacterial interference, that between pneumococci and *Staphylococcus aureus*, which, although well documented, occurs by an entirely unknown mechanism.

Several epidemiological studies have shown a negative association between carriage of *Streptococcus pneumoniae* and *S. aureus* (2, 3), raising public health concern that mass pneumococcal vaccination may cause an increase in *S. aureus* colonization and infection. As a case in point, it has been reported that children with recurrent otitis media vaccinated with the heptavalent pneumococcal vaccine had increased incidence of *S. aureus*-related acute otitis media and *S. aureus* colonization (3).

Recent in vitro and in vivo studies have demonstrated that the interference between these 2 pathogens is related to hydrogen peroxide production by *S. pneumoniae*, which is bactericidal to *S. aureus* (4, 5). Similar observations have been reported for certain other pairs of bacteria (6). It is highly intriguing how the relatively low levels of hydrogen peroxide produced safely by some bacteria are bactericidal to others, despite the relative abundance of mechanisms protecting bacterial cells from oxidative damage, such as H₂O₂-inactivating enzymes and antioxidants (7) or DNA lesion repair systems (8).

Here, we shed light on the mechanism of interference between H₂O₂-producing bacteria and *S. aureus*. We present data supporting the idea that prophages may have a much greater role in bacterial ecology than has hitherto been suspected—namely, that killing of a target organism by “remote control” prophage induction may represent a major modality of directional bacterial interference. We show also that lysogenic staphylococci are much more sensitive to DNA-damaging antibiotics, such as fluoroquinolones, than nonlysogens, almost certainly for the same reason. Given the high prevalence of lysogeny, we can now predict that small, SOS-inducing molecules, produced in the environment at subinhibitory concentrations, may have strong selective value as effectors of directional interference.

Results

Hydrogen Peroxide Kills Only Lysogenic *S. aureus*. Several species of bacteria have H₂O₂-dependent bactericidal activity toward *S. aureus* (4, 9). However, the mechanism by which the relatively low levels of H₂O₂ produced by these organisms are bactericidal to *S. aureus* remains to be determined. One possibility is that H₂O₂ produced by one organism induces the SOS response in a competing (target) organism, lethally activating resident prophages in the latter. If so, staphylococcal lysogens but not nonlysogens should be sensitive to H₂O₂ and pneumococci should be insensitive, even though they are often or always lysogenic. Accordingly, we tested 8 strains of *S. aureus*, 6 lysogenic, 2 nonlysogenic (RN450 and V329), including a congenic pair in which one (RN10359) was an 80α lysogen of the other (RN450) and a strain producing the phage-carried PVL toxin (strain LUG855), lately implicated in serious staphylococcal infections [10; supporting information (SI) Table S1]. We used H₂O₂ at 0.5 mM, in the range ordinarily seen with pneumococcal cultures, and observed that all of the lysogenic strains were highly

Author contributions: J.R.P. designed research; L.S. and D.V. performed research; G.R.-Y., K.T., J.M.C., and I.L. contributed new reagents/analytic tools; L.S., D.V., I.L., R.P.N., and J.R.P. analyzed data; and G.R.-Y., K.T., R.P.N., and J.R.P. wrote the paper;

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: jpenades@ivia.es.

This article contains supporting information online at www.pnas.org/cgi/content/full/0809600106/DCSupplemental.

© 2009 by The National Academy of Sciences of the USA

effects of DNA-damaging antibiotics than nonlysogens, owing to lysis caused by SOS-induced prophages in the former, as we have demonstrated above for H₂O₂. As shown in Fig. 4, two different lysogenic derivatives of *S. aureus* strain RN450, RN451 and RN10359, lysogenic for ϕ 11 and 80 α , respectively, were considerably more sensitive to ciprofloxacin (Fig. 4A) than their nonlysogenic parent and this sensitivity was eliminated in both cases by the phage repressor cleavage defect. Similarly, the noncleavable repressor mutation significantly reduced sensitivity to enrofloxacin (Fig. 4B). In a more general sense, we would predict that lysogenic bacteria would be more sensitive than nonlysogens to interference by antibiotic producers in the environment, where the antibiotic concentrations would surely be very low. This remains to be tested.

Materials and Methods

Bacterial Strains and Growth Conditions. Bacterial strains used in these studies are listed in Table S1. Bacteria were grown at 37 °C overnight on TSA agar medium (*S. aureus*) or BHI with 1.5% agar (*S. pneumoniae*), supplemented with antibiotics as appropriate. Broth cultures were grown at 37 °C in TSB (*S. aureus*) or Todd Hewitt (*S. pneumoniae*).

Induction of Prophages. Procedures for preparation and analysis of phage lysates, transduction, and transformation in *S. aureus* were performed essentially as described in refs. 17 and 18. In general, bacteria were grown in TSB to

OD₅₄₀ = 0.15 and induced by the addition of H₂O₂ at various concentrations and cultures were continued at 32 °C with slow shaking (80 rpm). Presence of the lysis was evaluated within 3 h by using previously applied criteria (17).

Coculture Assay. Interference between *S. aureus* and *S. pneumoniae* strains was measured in coculture assay as described in ref. 4. In brief, 1 mL of the *S. pneumoniae* cultures (OD₅₄₀ 0.4–0.5) were mixed with 1 mL of the *S. aureus* cultures (OD₅₄₀ 0.3–0.4), and incubated 4 h at 37 °C. To quantify bactericidal activity, the cocultured bacterial strains were plated on selective medium (TSA supplemented with 5 μ g/mL optochin), where only *S. aureus* can grow.

DNA Methods. General DNA manipulations were performed by standard procedures (19, 20). Strain JP3592 was obtained by using plasmid pMAD, as described in ref. 15. The oligonucleotides pairs used were ϕ 11–1cB (5'-cgcgatccAGTGT-TAATGTGTATATGCTC-3')/ ϕ 11–3c (5'-TAATTCTTCCTATCTCAGCACCAGTT-GCACC-3') and ϕ 80 α –cl-9mE (5'-CCGGAATTCACAGATTCGTTTATTTCCC-3')/ ϕ 11–4m (5'-GGTGCACTGGTGCTGAGATAGGAGAAGAATTA-3').

ACKNOWLEDGMENTS. We thank Simon Foster, Gabriela Bowden, and Pedro García for providing bacterial strains used in this study and Jeff Weiser, Simon Foster, Adela González de la Campa, and Alex Mira for helpful comments on the manuscript. This work was supported by Comisión Interministerial de Ciencia y Tecnología Grants BIO2005-08399-C02-02, BIO2008-05284-C02-02, and BIO2008-00642-E/C; Cardenal Herrera-CEU University Grants PRCEU-UCH25/08 and Copernicus program; and by Conselleria de Agricultura, Pesca i Alimentació (CAPiA), and from the Generalitat Valenciana (ACOMP07/258) (J.R.P.). L.S. and D.V. were supported by Cardenal Herrera-CEU University fellowships.

- Bassler BL, Losick R (2006) Bacterially speaking. *Cell* 125:237–246.
- Regev-Yochay G, et al. (2004) Association between carriage of *Streptococcus pneumoniae* and *Staphylococcus aureus* in Children. *JAMA* 292:716–720.
- Bogaert D, et al. (2004) Colonisation by *Streptococcus pneumoniae* and *Staphylococcus aureus* in healthy children. *Lancet* 363:1871–1872.
- Regev-Yochay G, Trzcinski K, Thompson CM, Malley R, Lipsitch M (2006) Interference between *Streptococcus pneumoniae* and *Staphylococcus aureus*: In vitro hydrogen peroxide-mediated killing by *Streptococcus pneumoniae*. *J Bacteriol* 188:4996–5001.
- Park B, Nizet V, Liu GY (2008) Role of *Staphylococcus aureus* catalase in niche competition against *Streptococcus pneumoniae*. *J Bacteriol* 190:2275–2278.
- Pericone CD, Overweg K, Hermans PW, Weiser JN (2000) Inhibitory and bactericidal effects of hydrogen peroxide production by *Streptococcus pneumoniae* on other inhabitants of the upper respiratory tract. *Infect Immun* 68:3990–3997.
- Cosgrove K, et al. (2007) Catalase (KatA) and alkyl hydroperoxide reductase (AhpC) have compensatory roles in peroxide stress resistance and are required for survival, persistence, and nasal colonization in *Staphylococcus aureus*. *J Bacteriol* 189:1025–1035.
- Goerlich O, Quillardet P, Hofnung M (1989) Induction of the SOS response by hydrogen peroxide in various *Escherichia coli* mutants with altered protection against oxidative DNA damage. *J Bacteriol* 171:6141–6147.
- Strauss E (2000) Microbiology. Fighting bacterial fire with bacterial fire. *Science* 290:2231–2233.
- Labandeira-Rey M, et al. (2007) *Staphylococcus aureus* Pantón-Valentine leukocidin causes necrotizing pneumonia. *Science* 315:1130–1133.
- Pericone CD, Park S, Imlay JA, Weiser JN (2003) Factors contributing to hydrogen peroxide resistance in *Streptococcus pneumoniae* include pyruvate oxidase (SpxB) and avoidance of the toxic effects of the fenton reaction. *J Bacteriol* 185:6815–6825.
- Novick RP, Subedi A (2007) The SaPIs: Mobile pathogenicity islands of *Staphylococcus*. *Chem Immunol Allergy* 93:42–57.
- Imlay JA, Chin SM, Linn S (1988) Toxic DNA damage by hydrogen peroxide through the Fenton reaction in vivo and in vitro. *Science* 240:640–642.
- Zhang X, et al. (2000) Quinolone antibiotics induce Shiga toxin-encoding bacteriophages, toxin production, and death in mice. *J Infect Dis* 181:664–670.
- Maiques E, et al. (2006) (beta)-Lactam antibiotics induce the SOS response and horizontal transfer of virulence factors in *Staphylococcus aureus*. *J Bacteriol* 188:2726–2729.
- Ubeda C, et al. (2005) Antibiotic-induced SOS response promotes horizontal dissemination of pathogenicity island-encoded virulence factors in staphylococci. *Mol Microbiol* 56:836–844.
- Novick RP (1991) Genetic systems in staphylococci. *Methods Enzymol* 204:587–636.
- Lindsay JA, Ruzin A, Ross HF, Kurepina N, Novick RP (1998) The gene for toxic shock toxin is carried by a family of mobile pathogenicity islands in *Staphylococcus aureus*. *Mol Microbiol* 29:527–543.
- Ausubel FM, et al. (1990) *Current Protocols in Molecular Biology* (Wiley, New York).
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab Press, Cold Spring Harbor, NY).